

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 31/445 // 31/70		A1	(11) International Publication Number: WO 98/30219						
			(43) International Publication Date: 16 July 1998 (16.07.98)						
(21) International Application Number: PCT/US98/00031		Glycobiology Institute, Dept. of Biochemistry, South Parks Road, Oxford OX1 3QU (GB).							
(22) International Filing Date: 13 January 1998 (13.01.98)		(74) Agents: MEYER, Scott, J. et al.; G.D. Searle & Co., Corporate Patent Dept., P.O. Box 5110, Chicago, IL 60680-5110 (US).							
(30) Priority Data: 08/782,321 13 January 1997 (13.01.97) US		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, GW, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).							
(63) Related by Continuation (CON) or Continuation-in-Part (CIP) to Earlier Application US 08/782,321 (CON) Filed on 13 January 1997 (13.01.97)		Published With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.							
(71) Applicant (for all designated States except US): MONSANTO COMPANY [US/US]; 800 North Lindbergh Boulevard, St. Louis, MO 63167 (US).									
(72) Inventors; and (75) Inventors/Applicants (for US only): PLATT, Frances, M. [GB/GB]; University of Oxford, Glycobiology Institute, Dept. of Biochemistry, South Parks Road, Oxford OX1 3QU (GB). NEISES, Gabrielle, R. [US/US]; 13939 Meursault, Chesterfield, MO 63017 (US). DWEK, Raymond, A. [GB/GB]; University of Oxford, Glycobiology Institute, Dept. of Biochemistry, South Parks Road, Oxford OX1 3QU (GB). BUTTERS, Terry, D. [GB/GB]; University of Oxford,									
(54) Title: METHOD FOR TREATMENT OF CNS-INVOLVED LYSOSOMAL STORAGE DISEASES									
(57) Abstract									
<p>A method is disclosed for the <i>in vivo</i> treatment of patients having a lysosomal storage disease with a significant central nervous system (CNS) involvement. Said method comprises administration to said patient a small but storage-inhibitory effective amount of an N-alkyl derivative of a 1,5-iminosugar in which said alkyl group contains from about 2 to about 8 carbon atoms and said 1,5-iminosugar is 1,5-dideoxy-1,5-imino-D-glucitol, or 1,5-dideoxy-1,5-imino-D-galactitol, or an O-acylated pro-drug of said 1,5-iminosugar. In an illustrative example, CNS storage of GM2 ganglioside is inhibited in Tay-Sachs mice by administration of 1,5-(butylimino)-1,5-dideoxy-D-glucitol.</p>									
<table border="1"> <caption>GM2 (arbitrary units) Data</caption> <thead> <tr> <th>Treatment</th> <th>GM2 (arbitrary units)</th> </tr> </thead> <tbody> <tr> <td>Untreated</td> <td>~27</td> </tr> <tr> <td>+ NB-DNJ</td> <td>~12</td> </tr> </tbody> </table>				Treatment	GM2 (arbitrary units)	Untreated	~27	+ NB-DNJ	~12
Treatment	GM2 (arbitrary units)								
Untreated	~27								
+ NB-DNJ	~12								

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon	KR	Republic of Korea	PL	Poland		
CN	China	KZ	Kazakhstan	PT	Portugal		
CU	Cuba	LC	Saint Lucia	RO	Romania		
CZ	Czech Republic	LJ	Liechtenstein	RU	Russian Federation		
DE	Germany	LK	Sri Lanka	SD	Sudan		
DK	Denmark	LR	Liberia	SE	Sweden		
EE	Estonia			SG	Singapore		

**METHOD FOR TREATMENT OF
CNS-INVOLVED LYSOSOMAL STORAGE DISEASES**

CROSS-REFERENCE TO RELATED APPLICATIONS

This is a continuation-in-part of application Ser. No.
08/393,640, filed February 24, 1995,

- which is a continuation of application Ser. No.
08/061,645, filed May 13, 1993, now U.S. Patent
5,399,567,
- and is a continuation-in-part of application Ser. No.
08/588,027, filed January 17, 1996,
- which is a division of application Ser. No. 08/396,989,
filed March 1, 1995,
- which is a division of application Ser. No. 08/102,654,
filed August 5, 1993,
- which is a continuation-in-part of said application
Ser. No. 08/061,645.

BACKGROUND AND FIELD OF THE INVENTION

This invention relates to a method for the treatment of lysosomal storage diseases that have a significant central nervous system (CNS) involvement. These diseases are caused by genetic mutations which result in the absence or deficiency of lysosomal enzymes. They include, for example, Tay-Sachs disease, Sandhoff disease, GM1 gangliosidosis and Fabry disease.

A list of references indicated by numerals in parentheses is appended at the end.

TAY-SACHS DISEASE:

This is a fatal hereditary disorder of lipid metabolism characterized especially in CNS tissue due to deficiency of the A (acidic) isozyme of β -hexosaminidase. Mutations in the *HEXA* gene, which encodes the α subunit of β -hexosaminidase, cause the A isozyme deficiency.

Tay-Sachs [disease] is a prototype of a group of disorders, the GM2 gangliosidoses, characterized by defective GM2 ganglioside degradation. The GM2 ganglioside (monosialylated ganglioside 2) accumulates in the neurons beginning already in fetal life.

SANDHOFF DISEASE:

Sandhoff disease results from a deficiency of both the A and B (basic) isozymes of β -hexosaminidase. Mutations in the *HEXB* gene, which encodes the β subunit of β -hexosaminidase, cause the B isozyme deficiency.

GM1 GANGLIOSIDOSIS:

GM1 gangliosidosis is caused by a deficiency of β -galactosidase, which results in lysosomal storage of GM1 ganglioside (monosialylated ganglioside 1).

FABRY DISEASE:

Fabry disease is caused by a deficiency of α -galactosidase which results in lysosomal storage of a ceramide trihexoside.

Glycosphingolipid (GSL) storage diseases are a group of human autosomal recessive disorders (except Fabry disease which is X-linked), each of which exhibits a characteristic pathology (1). They result from the inheritance of defects in genes encoding the catabolic enzymes required for the complete breakdown of GSLs within the lysosomes.

There presently is no effective therapy for Tay-Sachs disease or other lysosomal storage diseases with CNS involvement. Proposed strategies for the treatment of these debilitating and often fatal diseases include enzyme replacement therapy, gene therapy, substrate deprivation, allogenic bone marrow transplantation and palliative measures (2). Of these, symptomatic management is the only approach for treating most of these disorders, although transplantation techniques have been applied to some of these diseases.

Currently, only the non-neuronopathic form of Gaucher disease (type 1), a condition characterized by glucocerebrosidase deficiency, which occurs at high frequency in Ashkenazi Jews, has been successfully treated using enzyme replacement therapy (3,4). However, skeletal abnormalities associated with the disease respond slowly to this treatment (4) and the rare type 2 (acute neuronopathic; infantile) and rare type 3 (chronic; juvenile) are refractory to therapy.

Accordingly, new therapeutic treatment for lysosomal storage diseases which have significant CNS involvement (neuronopathic) are urgently needed.

BRIEF DESCRIPTION OF THE INVENTION

In accordance with the present invention, a method is provided for the in vivo treatment of patients having a lysosomal storage disease with a significant CNS involvement.

Said method comprises administration to said patient of a small but storage-inhibitory effective amount of an N-alkyl derivative of a 1,5-iminosugar in which said alkyl group contains from about 2 to about 8 carbon atoms and said 1,5-iminosugar is 1,5-dideoxy-1,5-imino-D-glucitol, or 1,5-dideoxy-1,5-imino-D-galactitol, or an O-acylated pro-drug of said 1,5-iminosugar.

Preferred 1,5-iminosugars are:

1,5-(Butylimino)-1,5-dideoxy-D-glucitol, which is also known as N-butyl deoxynojirimycin or by the abbreviated designation N-butyl DNJ;

1,5-(Butylimino)-1,5-dideoxy-D-galactitol, which is also known as N-butyl deoxygalactonojirimycin or by the abbreviated designation N-butyl DGJ;

1,5-(Butylimino)-1,5-dideoxy-D-glucitol, tetrabutyrate; and

1,5-(Butylimino)-1,5-dideoxy-D-galactitol, tetrabutyrate.

The method of the invention is illustrated in detail herein with the preferred compound, N-butyl DNJ. As described in detail herein, a mouse model of Tay-Sachs disease (9) is used to illustrate the in vivo effect of the 1,5-iminosugars for treatment of lysosomal storage diseases having a significant CNS involvement. Using this mouse model, it was demonstrated with the illustrative

N-butyl DNJ that this agent unexpectedly was able to cross the blood:brain barrier to an extent which inhibited CNS storage of GM2 ganglioside compared to the untreated control mice which exhibit progressive storage of that ganglioside.

DETAILED DESCRIPTION OF THE INVENTION

While the specification concludes with claims particularly pointing out and distinctly claiming the subject matter regarded as forming the invention, it is believed that the invention will be better understood from the following illustrative detailed description taken in conjunction with the accompanying drawings.

BRIEF DESCRIPTION OF THE DRAWINGS

FIG. 1 shows thin layer chromatography (TLC) analysis of GM2 ganglioside storage in the Tay-Sachs mouse in the presence or absence of N-butyl deoxynojirimycin (NB-DNJ). Mice were treated with NB-DNJ from 4 weeks of age up to twelve weeks and their GSL (glycosphingolipid) profiles compared at 4, 8 and 12 weeks relative to the untreated age matched controls. Each lane on the TLC plate represents the base-resistant GSLs derived from the whole brain of an individual mouse. The data are representative of studies carried out on five mice at each time point. The migration position of an authentic GM2 standard is indicated with an arrow.

FIG. 2, in two parts, A and B, shows prevention of GM2 storage in 12 week old mice. To demonstrate the variation in GM2 storage in mice treated with NB-DNJ, a group of three untreated and three NB-DNJ treated mice were compared at 12 weeks of age.

FIG. 2A: TLC profiles on total brain GSLs for three untreated mice (-) and three NB-DNJ treated mice (+).

FIG. 2B: Scanning densitometry on the GM2 species from FIG. 2A expressed in arbitrary units. The mean values +/- standard deviation are shown.

FIG. 3, in four parts, A-D, shows GM2 storage in the ventromedial hypothalamus of untreated and NB-DNJ treated mice (12 weeks of age). Frozen sections were stained with periodic acid-Schiff (PAS) to allow the visualization of GM2 storing neurons.

FIG. 3A: Untreated mouse, 10x magnification;

FIG. 3B: NB-DNJ treated mouse, 10x magnification;

FIG. 3C: Untreated mouse, 25x magnification; and

FIG. 3D: NB-DNJ treated mouse, 25x magnification.

The sections were selected to ensure that sections from the two animals were comparable in terms of spatial orientation within the brain. The images shown are representative of data derived from four different pairs of mice. The reduction in PAS staining in NB-DNJ treated mice was also observed in other storage regions of the brain.

FIG. 4, in four parts, A-D, shows electron microscopy of brains from untreated and NB-DNJ treated mice.

FIG. 4A: GM2 storage neuron from an untreated mouse brain;

FIG. 4B: GM2 storage neuron from an NB-DNJ treated mouse brain (the scale bar for A and B represents 1 μ m;

FIG. 4C: MCBs (membranous cytoplasmic bodies) from an untreated mouse brain;

FIG. 4D: MCBs from an NB-DNJ treated mouse brain (the scale bar for C and D represents 0.1 μ m).

The data shown are representative on the basis of analyzing multiple sections for multiple storage regions of the brain from two untreated and two NB-DNJ treated animals, with the analysis carried out by two independent groups.

FIG. 5 is a schematic flow chart which shows glycolipid catabolism and the relationship of lysosomal storage diseases to the absence or deficiency of relevant enzyme to degrade a given ganglioside or globoside. Of the lysosomal storage diseases that have a significant CNS involvement:

- * GM1 gangliosidosis is shown to be caused by a deficiency in β -galactosidase;
- * Tay-Sachs disease, which is a GM2 gangliosidosis, is shown to be caused by a deficiency of β -hexosaminidase A (acidic isozyme);
- * Sandhoff disease is shown to be caused by a deficiency of β -hexosaminidase A & B (acidic and basic isozymes); and
- * Fabry disease is shown to be caused by a deficiency in α -galactosidase.

In a different step of the GlcCer pathway, Gaucher disease is shown to be caused by a deficiency in β -glucocerebrosidase. Three diseases shown at the bottom of FIG. 5, namely metachromatic leukodystrophy (MLD), Krabbe disease and Niemann-Pick disease, involve different metabolic pathways, the GalCer pathway and sphingomyelinase (SM) pathway.

The mouse model of Tay-Sachs disease (9) is a useful model for demonstrating the in vivo effectiveness of the 1,5-iminosugars for the treatment of lysosomal storage diseases having a significant CNS involvement since this model has all the hallmarks of Tay-Sachs disease.

Tay-Sachs disease results from mutations in the HEXA gene, which encodes the α subunit of β -hexosaminidase, leading to a deficiency in the A isoenzyme. The A isoenzyme is responsible for the degradation of GM2 ganglioside. When this enzyme is deficient in humans, GM2 ganglioside accumulates progressively and leads to severe neurological degeneration (10).

In the mouse model of Tay-Sachs disease (generated by the targeted disruption of the Hexa gene), the mice store GM2 ganglioside in a progressive fashion, but the levels never exceed the threshold required to elicit neurodegeneration (9).

This is because in the mouse (but not human) a sialidase is sufficiently abundant that it can convert GM2 to GA2 (asialo ganglioside 2), which can then be catabolized by the hexosaminidase B isoenzyme (11). This model therefore has all the hallmarks of Tay-Sachs disease, in that it stores GM2 ganglioside in the CNS, but it never develops the neurological symptoms characteristic of the human disease (9,11,12).

In order to thus further illustrate the invention, the following detailed examples were carried out although it will be understood that the invention is not limited to these specific examples or the details described therein.

EXAMPLE I

Tay-Sachs mice were reared on standard mouse chow up to the age of weaning (4 weeks post-partum) when they were placed on a powdered mouse chow diet containing NB-DNJ as follows:

Mice were fed on a diet of powdered mouse chow (expanded Rat and Mouse Chow 1, ground, SDS Ltd., Witham, Essex, UK) containing NB-DNJ from weaning (4 weeks). The diet and compound (both dry solids) were mixed thoroughly before use, stored at room temperature and used within seven days of mixing. Water was available to the mice *ad lib*. The mice were housed under standard non-sterile conditions. The mice were given a dosing regime of 4800mg/kg/day of NB-DNJ which gave serum levels of approximately 50 μ M.

Similar serum levels (steady state trough level of approximately 20 μ M) were achieved in humans during the evaluation of this compound as an anti-viral agent when patients were treated with 43 mg/kg/day (12).

The pharmacokinetics of NB-DNJ are two orders of magnitude poorer in mouse relative to human, thereby necessitating high dosing regimes in the mouse in order to achieve serum levels in the predicted therapeutic range for the GSL storage disorders of 5-50 μ M (5-8).

EXAMPLE II

The effects that drug administration had on GM2 storage in the Tay-Sachs mouse were determined at various ages by extracting total brain lipids, separating the base resistant GSL fraction by TLC (FIG. 1), and identifying the GM2 species on the basis of co-migration with an authentic GM2 standard. The following procedure was employed:

The animals were anesthetized, perfused with phosphate buffered saline, pH 7.2, and the intact brain removed. The brain tissue was manually homogenized in water, freeze-dried and extracted twice with chloroform:methanol 2:1 (v/v) for two hours at room temperature and overnight at 4°C.

A volume of the solvent extract equivalent to 5 mg dry weight for each brain was dried under nitrogen, taken up in 500µl chloroform:methanol (1:1v/v), 83µl of 0.35M NaOH in 96% methanol added and incubated at room temperature for 90 mins. The samples were partitioned by adding 83µl H₂O:methanol (9:1 v/v), 166.5µl H₂O and 416µl chloroform, spun in a microfuge for 1 min and the upper phase retained. The lower phase was washed twice in Folch theoretical upper phase (chloroform:methanol:water, 1:10:10 v/v/v) and the upper phases retained and pooled with the original upper phase.

The samples were partially dried under N₂ to remove the solvent and the residual aqueous sample made up to 1ml with H₂O and dialyzed overnight in 2 liters of water to desalt. The samples were freeze-dried, extracted with 500µl chloroform:methanol 2:1 v/v, spun at 13000rpm for 2 mins and the supernatant retained, dried under N₂, resuspended in 10µl chloroform:methanol:0.22% calcium chloride (60:35:8 v/v/v) and separated by TLC (Silica gel

60 plates, Merck, BDH, Poole, Dorset, UK) in chloroform:methanol:calcium chloride (60:35:0.22%), sprayed with orcinol and visualized by heating to 80°C for 10 mins.

By four weeks of age a storage band corresponding to GM2 was detectable in the untreated mice, in agreement with previously published reports on this mouse model (9). As expected on the basis of published studies (9), the accumulation of GM2 in the untreated mice progressively increased with increasing age of the mice (FIG. 1).

However, in the NB-DNJ treated mice by eight weeks of age (four weeks untreated from birth to weaning, 4 weeks NB-DNJ treated post-weaning) there was an unexpected reduction in the intensity of the GM2 ganglioside band, relative to the untreated age matched controls, indicating that reduced levels of storage were occurring in the presence of the drug. The mice were followed for twelve weeks and there was a consistent reduction in stored GM2 ganglioside in all animals from the NB-DNJ treated group, irrespective of their age (FIG. 1).

To examine the generality of these data a group of three untreated and three NB-DNJ treated mice were evaluated at 12 weeks (FIG. 2A). In all cases, the intensity of the GM2 band was significantly reduced in the NB-DNJ treated animals, relative to the untreated age matched controls. When scanning densitometry was performed on the TLC profiles it was found that there was an approximately 50% reduction in GM2 ganglioside in the treated mouse brains relative to the untreated controls (FIG. 2B).

EXAMPLE III

The neurons within the Tay-Sachs mouse brains which are responsible for the GM2 storage observed in whole brain lipid extracts are confined to certain specific regions of the brain (12). We therefore carried out cytochemical analysis on tissue sections from untreated mice and mice treated for 16 weeks with NB-DNJ using periodic acid-Schiff (PAS) staining to detect the stored ganglioside within the storage neurons (9), as follows:

Mice were anesthetized, perfused with phosphate buffer pH 7.4 containing 4% paraformaldehyde and the brain dissected and retained in fixative overnight prior to cryopreservation and sectioning. Frozen brain sections (7 micron) were warmed to room temperature, stained with periodic acid-Schiff (PAS) according to the manufacturer's instructions (Sigma, Poole Dorset UK), counter-stained with Erhlich's hematoxylin and mounted in DPX (BDH).

It has previously been demonstrated in these untreated mice that the distribution of neurons staining with PAS is coincident with neurons which immunostain with an antibody specific for GM2 ganglioside (9).

In storage regions of the brain, such as the ventromedial hypothalamic nucleus, the NB-DNJ treated mice had fewer PAS positive neurons and the intensity of staining in each neuron was reduced (FIGS. 3B and 3D), relative to the untreated age matched control's brain sections, which exhibited extensive storage (FIGS. 3A and 3C).

The status of the GM2 storage in individual neurons from treated and untreated mouse brains was examined by electron microscopy (EM) as follows:

The mice were anesthetized and perfusion fixed with 2% paraformaldehyde, 2% glutaraldehyde mix in PBS. The brain was dissected and fixed in the same fixative overnight at 4°C. The brain was trimmed and 100µm sections cut on a vibratome, the sections washed three times in 0.1M phosphate buffer and stained with osmium tetroxide (1% in 0.1M phosphate) for 35 mins. The sections were dehydrated through an ethanol series, treated with propylene oxide (2 x 15 mins) and placed in Durcupan resin overnight at room temperature, transferred to glass slides and placed at 60°C for 48 hours.

Storage areas of the brain were selected microscopically, cut out of the thick section with a scalpel blade and glued with Super Glue Loctite, Quick Tite, (Loctite Corp., Rock Hill, CT) onto an Embed 800 stub (Electron Microscopy Sciences, Fort Washington, PA). Sections were stained with uranyl acetate/lead citrate and observed with a Hitachi 600 microscope at 75 kv.

In the storage neurons from untreated Tay-Sachs mouse brains there were prominent regions of the cytoplasm containing large numbers of membranous cytoplasmic bodies (MCBs) containing the stored lipid product (FIG. 4A). In contrast, in the NB-DNJ treated mice it proved difficult to find storage neurons. However, when storage cells could be located they contained MCBs with greatly reduced electron dense contents (FIG. 4B).

Furthermore, the extensive storage observed within storage neurons from untreated mice resulted in the organelles with the highest degree of storage being difficult to section, with the storage product frequently detaching partially from the surrounding membrane (FIG. 4A). In the NB-DNJ treated mouse brains the storage within neurons was always markedly reduced, relative to the untreated controls, and as a result no sectioning artifact was observed (FIG. 4B). This was consistently observed by two independent electron microscopy groups studying independent

material derived from these mice. One representative set of data is shown in FIG. 4.

The EM data are in keeping with the cytochemical staining which indicated that there were fewer storage neurons in the brains of treated mice and that storage cells in the treated animals had reduced levels of GM2 storage, relative to the untreated controls. When the morphology of individual MCBs from untreated and NB-DNJ treated mice were compared under high magnification by EM there was a profound difference in their morphology.

The NB-DNJ treated mice had MCBs which contained less electron dense storage lipid (FIG. 4D) but also did not have the prominent concentrically arranged lamellae characteristic of the MCB in neurons from untreated mice (FIG. 4C). Instead, they exhibited a diffuse pattern of storage with membrane-like structures only clearly discernible in the periphery of the organelle (FIG. 4D). Taken together with the cytochemical data, this demonstrates that NB-DNJ prevents lysosomal storage and the extent of storage per cell and per MCB is dramatically reduced, in keeping with the biochemical data on whole brain GSLs (FIGS. 1 and 2).

The data outlined herein demonstrate that oral treatment of mice with NB-DNJ is well tolerated and that it results in the inhibition of GSL biosynthesis. Furthermore, in the Tay-Sachs mouse, which exhibits progressive CNS storage of GM2 ganglioside, we have been able to prevent storage, as a consequence of reducing GSL biosynthesis. This indicates that NB-DNJ can cross the blood:brain barrier to an extent which can prevent storage.

This therefore indicates that substrate deprivation resulting from NB-DNJ administration is a rational strategy for the therapy of the human GSL lysosomal storage diseases. It has been shown in vitro that NB-DNJ specifically inhibits the first step in GSL

biosynthesis, the glucosyl-transferase catalyzed biosynthesis of GlcCer (3-7).

As several of the human glycosphingolipid (GSL) storage diseases involve the storage of GlcCer-based GSLs, this therapeutic strategy can be applied to all of these disease states, irrespective of the specific storage product. This would include Gaucher (types 1, 2 and 3), Fabry disease, Tay-Sachs disease, Sandhoff disease, GM1 gangliosidosis, and fucosidosis.

The current application of enzyme replacement to Gaucher disease is limited by the fact that the enzyme cannot cross the blood:brain barrier and hence this therapy is only efficacious in type 1 disease where there is no neuropathology involved. Our finding that GSL depletion can be achieved in the central nervous system is therefore of major significance as it means that all the GSL storage diseases could be treated with NB-DNJ, as many of them involve neuropathology in the CNS.

NB-DNJ does not appear to inhibit galactosyltransferase which initiates the biosynthesis of the pathway that results in the formation of GalCer and sulphatide. Therefore, it is not believed that NB-DNJ would show efficacy against Krabbe disease and metachromatic leukodystrophy (MLD). These diseases involve the storage of GalCer and sulphatide, respectively, as shown at the bottom of FIG. 5. This can be advantageous to the invention as the formation of GalCer and sulphatide, which are important constituents of myelin, would not be affected by the treatment. Hence, myelination and myelin stability would not be impaired.

EXAMPLE IV

When any of the following compounds are substituted for an equivalent amount of 1,5-(Butylimino)-1,5-dideoxy-D-glucitol in the above Examples I, II and III, substantially similar inhibitory results are obtained:

- A) 1,5-(Hexylimino)-1,5-dideoxy-D-glucitol;
- B) 1,5-(Butylimino)-1,5-dideoxy-D-galactitol;
- C) 1,5-(Hexylimino)-1,5-dideoxy-D-galactitol;
- D) 1,5-(Butylimino)-1,5-dideoxy-D-glucitol, tetrabutyrates;
- E) 1,5-(Hexylimino)-1,5-dideoxy-D-glucitol, tetraacetate.

Compounds D and E are synthesized as described in U.S. Patent 5,003,072.

In treatment of the recipient patients in accordance with the method of the invention, the active agent can be administered by conventional drug administration procedures, preferably in formulations with pharmaceutically acceptable diluents and carriers. The active agent can be used in the free amine form or in the salt form. Pharmaceutically acceptable salt forms are illustrated, for example, by the HCl salt.

The amount of active agent to be administered must be an effective amount, that is, an amount which will be medically beneficial but does not present toxic effects which outweigh the advantages which accompany its use. It would be expected that the

average adult human daily dosage would normally range from about 0.1 mg to about 1000 mg of the active agent.

The preferable route of administration is orally in the form of capsules, tablets, syrups, elixirs and the like, although parenteral administration can also be had. Suitable formulations of the active compound in pharmaceutically acceptable diluents and carriers in therapeutic dosage form can be prepared by conventional procedures such as by reference to general texts in the field, e.g., Remington's Pharmaceutical Sciences, ed. Arthur Osol, 16th ed., 1980, Mack Publishing Co., Easton, PA, and the 18th ed., 1990. Conventional diluents and carriers are, e.g., water, normal saline, sugars, starch and the like substances.

Various other examples will be apparent to the person skilled in the art after reading the disclosure herein. All such other examples are meant to be included within the scope of the appended claims.


REFERENCES

1. Neufeld, E.F. (1991) *Ann. Rev. Biochem.*, 60, 257-280.
2. Beutler, E. (1992) *Science*, 256, 794-799.
3. Barton, N.W., Brady, R.O., Dambrosia, J.M., Di Bisceglie, A.M., Doppelt, S.H., Hill, S.C., Mankin, H.J., Murray, G.J., Parker, R.I., Argoff, C.E., Grewal, R.P., Yu, K-T., and Collaborators (1991) *N. Eng. J. Med.* 324, 1464-1470.
4. Beutler, E., Kay, A., Saven, A., Garver, P., Thurston, D., Dawson, A. and Rosenbloom, B. (1991) *Blood*, 78, 1183-1189.
5. Platt, F.M., Neises, G. R., Dwek, R.A. and Butters, T.D. (1994) *J. Biol Chem.*, 269, 8362-8365.
6. Platt, F.M., Neises, G. R., Dwek, R.A. and Butters, T.D. (1994) *J. Biol Chem.*, 269, 27108-27114.
7. Platt, F.M. and Butters, T.D. (1995) Inhibitors of Glycosphingolipid Biosynthesis, *Trends in Glycoscience and Glycotechnology*. 269, 495-511.
8. Yamanaka, S., Johnson, M.D., Grinberg, A., Westphal, H., Crawley, J.N., Taniike, M., Suzuki, K. and Proia, R.L. (1994) *Proc. Natl. Acad. Sci. U.S.A.*, 91, 9975-9979.
9. Sandhoff, K., Conzelmann, E., Neufeld, E.F., Kaback, M.M., and Suzuki, K. (1989) in *The Metabolic Basis of Inherited Disease*, eds. Scriver, C.R., Beaudet, A.L., Sly, W.S. and Valle, D. (McGraw-Hill, New York, Vol. 2, pp1807-1839.
10. Sango, K., Yamanaka, S. Hoffmann, A., Okuda, Y., Grinberg, A., Westphal, H., McDonald, M.P., Crawley, J.N., Sandhoff, K., Suzuki, K. and Proia, R.L. (1995) *Nature Genet.* 11, 170-176.
11. Taniike, M., Yamanaka, S., Proia, R.L., Langaman, C., Bone-Turrentine, T. and Suzuki, K. (1995) *Acta Neuropathol.* 89, 296-304.
12. Fischl, M.A., Resnick, L., Coombs, R., Kremer, A.B., Pottage, J.C., Pass, R.J., Pife, K.H., Powderly, W.G., Collier, A.C., Aspinall, R.L., Smith, S.L., Kowalski, K.G., and Wallemark, C-B. (1994) *J. AIDS* 7, 139-147.

WHAT IS CLAIMED:


1. A method for the treatment of patients having a lysosomal storage disease with a significant central nervous system (CNS) involvement comprising administering to said patient a small but storage-inhibitory effective amount of an N-alkyl derivative of a 1,5-iminosugar in which said alkyl group contains from about 2 to about 8 carbon atoms and said 1,5-iminosugar is a 1,5-dideoxy-1,5-imino-D-glucitol, or 1,5-dideoxy-1,5-imino-D-galactitol or an O-acylated prodrug of said 1,5-iminosugar.
2. The method of Claim 1 in which the 1,5-iminosugar is 1,5-dideoxy-1,5-imino-D-glucitol.
3. The method of Claim 1 in which the alkyl group is butyl.
4. The method of Claim 1 in which the 1,5-iminosugar is 1,5-dideoxy-1,5-imino-D-glucitol and the alkyl group is butyl.
5. The method of Claim 4 in which the 1,5-iminosugar is O-acylated to form the tetrabutyrates.
6. The method of any of Claims 1-5 in which the lysosomal storage disease is Tay-Sachs disease.

1/6

*FIG. 1*

2/6


NB-DNJ



← GM2

FIG. 2A

3 / 6

*FIG. 2B*

4/6

FIG. 3A




FIG. 3B




FIG. 3C



FIG. 3D

FIG. 4A



FIG. 4B



FIG. 4C




FIG. 4D

6 / 6

Glycolipid Catabolism




FIG. 5

INTERNATIONAL SEARCH REPORT

national Application No

PCT/US 98/00031

A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 A61K31/445 //A61K31/70

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 6 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the International search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	PLATT ET AL.: "Prevention of lysosomal storage in Tay-Sachs mice treated with N-butyldeoxynojirimycin" SCIENCE, vol. 276, no. 5311, 18 April 1997, pages 428-431, XP002065772 see the whole document --- -/--	1-4,6

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

25 May 1998

Date of mailing of the international search report

11.06.98

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 opo nl,
Fax: (+31-70) 340-3016

Authorized officer

Gac, G

INTERNATIONAL SEARCH REPORT

International Application No.

PCT/US 98/00031

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	PLATT ET AL.: "Extensive glycosphingolipid depletion in the liver and lymphoid organs of mice treated with N-butyldeoxynojirimycin" J. BIOL. CHEM., vol. 272, no. 31, 1 August 1997, pages 19365-19372, XP002065773 see page 19365, right-hand column, paragraph 3 see page 18371, left-hand column, paragraph 2 see page 19372, left-hand column, paragraph 2 ---	1-6
P,X	KOLTER ET AL.: "A chemical concept for the treatment of Tay-Sachs disease" ANGEW. CHE., INT. ED. ENGLAND, vol. 36, no. 18, 2 October 1997, pages 1955-1959, XP002065774 see the whole document ---	1-4,6
X	WO 94 26714 A (G.D. SEARLE & CO.) 24 November 1994 cited in the application see the whole document, especially pages 2-4, 16-21, and page 25 lines 16-21 ---	1-5
Y	---	6
X	DANIEL ET AL.: "Evidence for processing of dolichol-linked oligosaccharides in patients with neuronal ceroid-lipofuscinosis" AM. J. MED. GENET., vol. 42, no. 4, 15 February 1992, pages 586-592, XP002065775 see page 586 see page 590, right-hand column, last paragraph ---	1,2
Y	LIESSEM ET AL.: "Synthese von 2-acetamido-1,4-imino-1,2,4-tridesoxy-D-Ga laktitol und kompetitive Inhibition der humanen lysosomalen beta-Hexosaminidase A" CARBOHYDRATE RES., vol. 250, no. 1, 1993, pages 19-30, XP002065776 see page 20 ---	6
A	---	1-5

	-/--	

INTERNATIONAL SEARCH REPORT

national Application No
PCT/US 98/00031

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	PLATT ET AL.: "N-butyldeoxygalactonojirimycin inhibits glycolipid biosynthesis but does not affect N-linked oligosaccharide processing" J. BIOL. CHEM., vol. 268, no. 43, 28 October 1994, pages 27108-27115, XP002065777 see the whole document ---	1-6
A	PLATT F M ET AL: "N-BUTYLDEOXYNOJIRIMYCIN IS A NOVEL INHIBITOR OF GLYCOLIPID BIOSYNTHESIS" JOURNAL OF BIOLOGICAL CHEMISTRY, vol. 269, no. 11, 18 March 1994, pages 8362-8365, XP000615445 ---	1-4,6
A	OSIECKI-NEWMAN K ET AL: "HUMAN ACID -GLUCOSIDASE: USE OF INHIBITORS ALTERNATIVE SUBSTRATES AND AMPHIPHILES TO INVESTIGATE THE PROPERTIES OF THE NORMAL AND GAUCHER DISEASE ACTIVE SITES" BIOCHIMICA ET BIOPHYSICA ACTA, vol. 915, no. 1, 1 January 1987, pages 87-100, XP000561731 ---	1-4
A	LEMBCKE ET AL.: "Lysosomal storage of glycogen as a sequel of alpha-glucosidase inhibition by the absorbed deoxynojirimycin derivative emiglitate (BAYo1248)" RES. EXP. MED., vol. 191, no. 6, 1991, BERLIN, pages 389-404, XP002065778 see the whole document -----	1

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US 98/00031

Box I Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

Although claims 1-6 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of Item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Information on patent family members

national Application No

PCT/US 98/00031

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9426714 A	24-11-1994	US 5399567 A	21-03-1995
		AT 148456 T	15-02-1997
		AU 6783294 A	12-12-1994
		CA 2159988 A	24-11-1994
		DE 69401658 D	13-03-1997
		DE 69401658 T	12-06-1997
		EP 0698012 A	28-02-1996
		ES 2097653 T	01-04-1997
		JP 8510244 T	29-10-1996
		US 5472969 A	05-12-1995
		US 5656641 A	12-08-1997
		US 5580884 A	03-12-1996
		US 5525616 A	11-06-1996